

## DISCLAIMER

All labeling reflected on this website is for informational and promotional purposes only. It is not intended to be used by healthcare professionals or patients for the purpose of prescribing or administering these products. Questions regarding the current content of product labeling should be directed to Akom's Customer Service department at 800.932.5676.

# Albuterol Sulfate Inhalation Solution, 0.5%\* Sterile

(\*Potency expressed as albuterol)

## Patient's Instructions for Use

**Note: The albuterol sulfate inhalation solution contained in the 20 mL multiple-dose bottle is concentrated and must be diluted.**

**Read complete instructions carefully before using.**

1. Draw the appropriate volume of albuterol sulfate inhalation solution, 0.5% into the specially marked dropper that comes with each multidose bottle (Figure 1). For children 12 years of age and under, the volume is based upon body weight. Use the dropper volume prescribed by your doctor.



Figure 1

2. Squeeze the solution into the nebulizer reservoir through the appropriate opening, taking care not to touch the tip of the dropper (Figure 2).




Figure 2

3. Add sterile normal saline solution, as your doctor has directed. A general guideline for the amount of saline to add is: For children using 0.25 mL or 1.25 mg of albuterol sulfate inhalation solution, add 2.75 mL of sterile normal saline. For children or adults using 0.5 mL or 2.5 mg of albuterol sulfate inhalation solution, add 2.5 mL of sterile normal saline.

4. Gently swirl the nebulizer to mix the contents and connect it with the mouthpiece or face mask (Figure 3).



Figure 3

5. Connect the nebulizer to the compressor.

6. Sit in a comfortable, upright position; place the mouthpiece in your mouth (Figure 4) (or put on the face mask); and turn on the compressor.



Figure 4

7. **Breathe as calmly, deeply, and evenly** as possible until no more mist is formed in the nebulizer chamber (about 5 to 15 minutes). At this point, the treatment is finished.
8. Clean the nebulizer (see manufacturer's instructions).

**Note: Use only as directed by your physician. More frequent administration or higher doses are not recommended.**

**Store albuterol sulfate inhalation solution, 0.5%\* between 2° and 25°C (36°-77°F).**

To avoid microbial contamination, proper aseptic techniques should be used each time the bottle is opened. Precautions should be taken to prevent contact of the dropper tip of the bottle with any surface, including the nebulizer reservoir and associated ventilatory equipment. In addition, if the solution changes color or becomes cloudy, it should not be used. The safety and effectiveness of albuterol sulfate inhalation solution have not been determined when one or more drugs are mixed with it in a nebulizer. Check with your doctor before mixing any medications in your nebulizer.

Call your doctor for medical advice about side effects. You may report side effects to Hi-Tech Pharmacal Co., Inc. at 1-800-262-9010 or FDA at 1-800-FDA-1088.

ADDITIONAL INSTRUCTIONS:

Hi-Tech Pharmacal Co., Inc.  
Amityville, NY 11701  
PI191 Rev. 01/18  
MG034 Rev. 10/18