


DISCLAIMER

All labeling reflected on this website is for informational and promotional purposes only. It is not intended to be used by healthcare professionals or patients for the purpose of prescribing or administering these products. Questions regarding the current content of product labeling should be directed to Akom's Customer Service department at 800.932.5676.

Tobramycin Ophthalmic Solution, USP 0.3%—Sterile



GTM00N


Rev. 03/13

R_x only

DESCRIPTION:

Tobramycin Ophthalmic Solution USP, 0.3% is a sterile topical ophthalmic antibiotic formulation prepared specifically for topical therapy of external ophthalmic infections.

Tobramycin is a water soluble aminoglycoside antibiotic active against a wide variety of gram-negative and gram-positive ophthalmic pathogens. The molecular formula is $C_{18}H_{37}N_5O_9$, the molecular weight is 467.52 and the structural formula is:



Chemical Name:

O-[3-amino-3-deoxy- α -D-glucopyranosyl-(1 \rightarrow 4)]-O-[2,6-diamino-2,3,6-trideoxy- α -D-ribohexo-pyranosyl-(1 \rightarrow 6)]-2-deoxystreptamine.

Each mL contains:

Active: Tobramycin 3 mg (0.3%). **Inactives:** Boric Acid, Sodium Chloride, Sodium Sulfate, Tyloxapol, Sodium Hydroxide and/or Sulfuric Acid to adjust pH (7.0 to 8.0), and Purified Water USP. **Preservative:** Benzalkonium Chloride 0.1 mg (0.01%).

CLINICAL PHARMACOLOGY:

In Vitro Data: *In Vitro* studies have demonstrated tobramycin is active against susceptible strains of the following microorganisms:

Staphylococci, including *S. aureus* and *S. epidermidis* (coagulase-positive and coagulase-negative), including penicillin-resistant strains.

Streptococci, including some of the Group A beta-hemolytic species, some nonhemolytic species, and some *Streptococcus pneumoniae*.

Pseudomonas aeruginosa, *Escherichia coli*, *Klebsiella pneumoniae*, *Enterobacter aerogenes*, *Proteus mirabilis*, *Morganella morganii*, most *Proteus vulgaris* strains, *Haemophilus influenzae* and *H. aegyptius*, *Moraxella lacunata*, *Acinetobacter calcoaceticus* and some *Neisseria* species.

Bacterial susceptibility studies demonstrate that in some cases, microorganisms resistant to gentamicin retain susceptibility to tobramycin.

INDICATIONS AND USAGE:

Tobramycin Ophthalmic Solution USP, 0.3% is a topical antibiotic indicated in the treatment of external infections of the eye and its adnexa caused by susceptible bacteria. Appropriate monitoring of bacterial response to topical antibiotic therapy should accompany the use of Tobramycin Ophthalmic Solution USP, 0.3%. Clinical studies have shown tobramycin to be safe and effective for use in pediatric patients.

CONTRAINDICATIONS:

Tobramycin Ophthalmic Solution USP, 0.3% is contraindicated in patients with known hypersensitivity to any of its components.

WARNINGS:

FOR TOPICAL OPHTHALMIC USE ONLY. NOT FOR INJECTION INTO THE EYE. Sensitivity to topically applied aminoglycosides may occur in some patients. If a sensitivity reaction to Tobramycin Ophthalmic Solution USP, 0.3% occurs, discontinue use.

PRECAUTIONS:

General:

As with other antibiotic preparations, prolonged use may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Cross-sensitivity to other aminoglycoside antibiotics may occur; if hypersensitivity develops with this product, discontinue use and institute appropriate therapy. Patients should be advised not to wear contact lenses if they have signs and symptoms of bacterial conjunctivitis.

Information for Patients: Do not touch dropper tip to any surface, as this may contaminate the solution.

Pregnancy Category B: Reproduction studies in three types of animals at doses up to thirty-three times the normal human systemic dose have revealed no evidence of impaired fertility or harm to the fetus due to tobramycin. There are, however, no adequate and well-controlled studies in pregnant women. Because animal studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers: Because of the potential for adverse reactions in nursing infants from tobramycin, a decision should be made whether to discontinue nursing the infant or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use: Safety and effectiveness in pediatric patients below the age of two months has not been established.

Geriatric Use: No overall clinical differences in safety or effectiveness have been observed between the elderly and other adult patients.

ADVERSE REACTIONS:

The most frequent adverse reactions to tobramycin ophthalmic solution are hypersensitivity and localized ocular toxicity, including lid itching and swelling, and conjunctival erythema. These reactions occur in less than three of 100 patients treated with tobramycin. Similar reactions may occur with the topical use of other aminoglycoside antibiotics. Other adverse reactions have not been reported from tobramycin therapy; however, if topical ocular tobramycin is administered concomitantly with systemic aminoglycoside antibiotics, care should be taken to monitor the total serum concentration.

OVERDOSAGE:

Clinically apparent signs and symptoms of an overdose of tobramycin ophthalmic solution (punctate keratitis,

erythema, increased lacrimation, edema and lid itching) may be similar to adverse reaction effects seen in some patients.

DOSAGE AND ADMINISTRATION:

In mild to moderate disease, instill one or two drops into the affected eye(s) every four hours. In severe infections, instill two drops into the eye(s) hourly until improvement, following which treatment should be reduced prior to discontinuation.

HOW SUPPLIED:

Tobramycin Ophthalmic Solution USP, 0.3% is supplied as a sterile solution in plastic dropper bottles in two sizes:

NDC 17478-290-20 2 mL

NDC 17478-290-10 5 mL

STORAGE:

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep container tightly closed. Protect from excessive heat.



Manufactured by: **Akorn, Inc.**
Lake Forest, IL 60045