

DISCLAIMER

All labeling reflected on this website is for informational and promotional purposes only. It is not intended to be used by healthcare professionals or patients for the purpose of prescribing or administering these products. Questions regarding the current content of product labeling should be directed to Akorn's Customer Service department at 800.932.5676.

Guaiatussin AC

Sugar-Free

Expectorant/Cough Suppressant

Description: Each 5 mL (1 teaspoonful) contains:

Guaifenesin.....	100 mg
Codeine Phosphate.....	10 mg
Alcohol	3.5% v/v

Inactive ingredients:

Alcohol 3.5% v/v, artificial cherry flavor, caramel, citric acid, FD&C red # 40, glycerin, polyethylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution.

Uses • temporarily relieves cough due to minor throat and bronchial irritations as may occur with the common cold or inhaled irritants • helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive

Warnings

Do not use • in adults and children who have a chronic pulmonary disease or shortness of breath, or children who are taking other drugs, unless directed by a doctor.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives, tranquilizers and drugs used for depression, especially monoamine oxidase inhibitors (MAOIs). These combinations may cause greater sedation (drowsiness) than is caused by the product used alone.

Stop use and ask a doctor if • cough lasts more than 7 days, comes back, or occurs with fever, rash or headache that lasts. These can be signs of a serious condition.
• may cause or aggravate constipation

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. • Use of codeine-containing preparation is not recommended for children under 2 years of age

Directions • take every 4 hours • do not exceed 6 doses in 24 hours • a special measuring device should be used to give an accurate dose of this product to children under 6 years of age • giving a higher dose than recommended by a doctor can result in serious side effects for a child

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls (10 mL)
Children 6 years to under 12 years of age	1 teaspoonful (5 mL)
Children under 6 years of age	Consult a doctor

How supplied: Guaiatussin AC Sugar – Free is a light amber – colored, viscous liquid with a cherry odor supplied in the following oral dosage forms: 5 mL unit dose, 10 mL unit dose in trays of 10, 4 fl. oz. (118 mL), 8 fl. oz. (237 mL) and 16 fl. oz. (473 mL) bottles.

Each (5 mL) teaspoon contains: 1 mg sodium.

Storage: Keep container closed and away from heat. Store at room temperature 15°- 30°C (59°- 86°F).

Questions or comments? Call 1-800-262-9010.

Hi-Tech Pharmacal Co., Inc.
Amityville, NY 11701

Rev.087:00 2/12
MG #32163