All labeling reflected on this website is for informational and promotional purposes only. It is not intended to be

used by healthcare professionals or patients for the purpose of prescribing or administering these products.

Questions regarding the current content of product labeling should be directed to Akorn's Customer Service

DISCLAIMER

department at 800.932.5676.

Drug Facts

Active ingredients (in each 5mL teaspoonful) Purposes

Codeine Phosphate, USP 10 mg......Cough Suppressant Guaifenesin, USP 100 mg.....Expectorant

Use • temporarily relieves cough due to minor throat and bronchial irritations as may occur with the common cold or inhaled irritants

 helps loosen phlegm (mucus) and thin bronchial secretions to make cough more productive

Warnings

Do not use • in adults and children who have a chronic pulmonary disease or shortness of breath, or children who are taking other drugs, unless directed by a doctor.

Ask a doctor before use if you have • a cough with too much phlegm (mucus) • a persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives, tranquilizers and drugs used for depression, especially monoamine oxidase inhibitors (MAOIs). These combinations may cause greater

sedation (drowsiness) than is caused by the product used alone.

Stop use and ask a doctor if • cough lasts for more than 7 days, comes back, or occurs with fever, rash or headache that lasts. These can be signs of a serious condition. • may cause or aggravate constipation

for pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately. • Use of codeine-containing preparation is not recommended for children under 2 years of age



PHARMACAL

NDC 50383-087-16

Guaiatussin AC Sugar-Free

- Expectorant
- Cough Suppressant

Hi-Tech Pharmacal Co., Inc. Amityville, NY 11701

16 fl oz (473 mL)

Drug Facts (continued)

children under 6 years of age

Directions • take every 4 hours • do not exceed 6 doses in 24 hours • a special measuring device should be used to give an accurate dose of this product to children under 6 years of age • giving a higher dose than recommended by a doctor can result in serious side effects for a child adults and children 12 years and over 2 teaspoonful children 6 to under 12 years of age 1 teaspoonful

Consult a doctor

Other information • each (5 mL) teaspoon contains: 1 mg sodium • Tamper evident: Do not use if seal under cap is broken or missing • Keep container closed and store away from heat • Store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients alcohol 3.5% v/v, artificial cherry flavor, caramel, citric acid, FD&C red # 40, glycerin, polyethylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution.

Questions or comments?

Call **1-800-262-9010**, Mon. - Thurs. 9:00 am thru 4:30 pm EST, Fri. 9:00 am thru 2:30 pm EST.

Serious side effects associated with use of this product may be reported to this number.



50383-087-16

REV. 087:01 9/11