

Loperamide Hydrochloride Oral Solution

1 mg/5 mL

Anti-Diarrheal

Rev. 618:02 01/15



Drug Facts

Active ingredient

(in each 5 mL teaspoonful)

Purpose

Loperamide HCl 1 mgAntidiarrheal

Uses controls the symptoms of diarrhea, including Traveler's Diarrhea.

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Do not use if you have bloody or black stool.

Ask a doctor before use if you have

- high fever (greater than 101°F)
 - mucus present in your stool
 - a history of liver disease
-

Ask a doctor or pharmacist before use if you are taking antibiotics.

Stop use and ask a doctor if diarrhea lasts more than 2 days.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluid to help prevent dehydration, which may accompany diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

Drug Facts (continued)

adults and children 12 years and older	20 mL after the first loose bowel movement; 10 mL after each subsequent loose bowel movement; but no more than 40 mL a day
children 9-11 years (60-95 lbs)	10 mL after the first loose bowel movement; 5 mL after each subsequent loose bowel movement; but no more than 30 mL a day
children 6-8 years (48-59 lbs)	10 mL after the first loose bowel movement; 5 mL after each subsequent loose bowel movement; but no more than 20 mL a day
children under 6 years (up to 47 lbs)	ask a doctor (not intended for use in children under 6 years old)

Other information

- avoid excess heat
- store at room temperature, 15°C to 30°C (59°F to 86°F)

Inactive ingredients alcohol (0.5%), benzoic acid, cherry flavor, glycerin, mint flavor, propylene glycol, purified water, sodium benzoate, sorbitol solution. Citric acid may be used to adjust pH.

Questions or comments?

Call **1-800-262-9010** Mon. – Thurs. 9:00 am – 4:30 pm EST, Fri. 9:00 am thru 2:30 pm EST. Serious side effects associated with use of this product may be reported to this number.

Manufactured by:
Hi-Tech Pharmacal Co., Inc.
Amityville, NY 11701

Rev. 618:02 01/15
MG #36488