

DISCLAIMER

All labeling reflected on this website is for informational and promotional purposes only. It is not intended to be used by healthcare professionals or patients for the purpose of prescribing or administering these products. Questions regarding the current content of product labeling should be directed to Akorn's Customer Service department at 800.932.5676.

DOCU LIQUID
(Docusate Sodium)
STOOL SOFTENER LAXATIVE

Rev. 771:01 4/13



DESCRIPTION: Each 5 mL (teaspoonful) contains docusate sodium 50 mg. *Inactive Ingredients:* D&C Red No. 33, methylparaben, natural and artificial vanilla flavor, poloxamer 181, polyethylene glycol, propylene glycol, propylparaben, purified water, sodium benzoate. Sodium citrate may be used to adjust pH.

Sodium Content: 5 mg/5 mL.

INDICATIONS: For the relief of occasional constipation due to hard stools. The effect on stools is usually apparent 1 to 3 days after the first dose.

DIRECTIONS: As directed by a physician or once daily as follows: *Adults and children 12 years of age and over:* oral dosage is 5 to 20 mL (1 to 4 teaspoonfuls). *Children 6 to under 12 years of age:* oral dosage is 5 to 10 mL (1 to 2 teaspoonfuls). *Children 3 to under 6 years of age:* oral dosage is 2.5 to 5 mL (1/2 to 1 teaspoonful). *Children under 3 years of age:* consult a physician. The higher doses are recommended for initial therapy. Dosage should be adjusted to individual response. If desired, the product may be administered in a 6 to 8 oz glass of milk or fruit juice or in infant formula to mask the bitter taste.

WARNINGS: Do not use laxative products when abdominal pain, nausea, or vomiting are present, or use for a period longer than 1 week, unless directed by a physician. If you have noticed a sudden change in bowel habits that persists over a period of 2 weeks, consult a physician before using a laxative. Rectal bleeding or failure to have a bowel movement after the use of a laxative may indicate a serious condition. Discontinue use and consult your physician. May cause rash in some cases.

If pregnant or breast-feeding, ask a health professional before use.

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN. In case of accidental overdose, seek professional assistance or contact a Poison Control Center immediately.

DRUG INTERACTION PRECAUTION: Do not take this product if you are presently taking mineral oil or a prescription drug, unless directed by a physician.

HOW SUPPLIED: Pink colored, liquid supplied in the following oral dosage forms: unit dose cups of 10 mL, 10 cups per tray and 16 fl. oz. (473 mL) bottle.

STORAGE: Keep tightly closed. Store at controlled room temperature, 15°-30°C (59°-86°F). Protect from freezing. Protect from light.

Questions or comments? Call **1-800-262-9010**

Hi-Tech Pharmacal Co., Inc.

Amityville, NY 11701

Rev. 771:01 4/13

MG #28534